

A MODEL-FREE VERSION OF THE FUNDAMENTAL THEOREM OF ASSET PRICING AND THE SUPER-REPLICATION THEOREM

B. ACCIAIO^{†‡}, M. BEIGLBÖCK[‡], F. PENKNER[‡], AND W. SCHACHERMAYER[‡]

ABSTRACT. We propose a *Fundamental Theorem of Asset Pricing* and a *Super-Replication Theorem* in a model-independent framework. We prove these theorems in the setting of finite, discrete time and a market consisting of a risky asset S as well as options written on this risky asset. As a technical condition, we assume the existence of a traded option with a super-linearly growing payoff-function, e.g., a power option. This condition is not needed when sufficiently many vanilla options maturing at the horizon T are traded in the market.

Keywords: Model-independent pricing, Fundamental Theorem of Asset Pricing, Super-Replication Theorem.
Mathematics Subject Classification (2010): 91G20, 60G42

1. INTRODUCTION

We consider a finite, discrete time setting and a market consisting of a collection of options $\varphi_i, i \in I$ written on a risky asset S . We work in complete generality, allowing I to be any set and the φ_i any kind of (possibly path-dependent) options written on S . In this context we address the following questions:

- (Q1) Does there exist an arbitrage opportunity?
- (Q2) For any additional option written on S , what is the range of prices that do not create an arbitrage opportunity?

These questions have been widely investigated and exhaustively answered in the classical model-dependent framework, where assumptions are made on the dynamics of the underlying process S , see [Sch10, Cam10] and the references therein.

We study these problems without making any model assumption. Instead, we consider the set of all models which are compatible with the prices observed in the market, i.e., we follow the *model-independent approach* to financial mathematics. A particular case is the situation when one observes the prices of finitely many European call options. This is the setup studied in Davis and Hobson [DH07], where the authors identify three possible cases: *absence of arbitrage*, *model-independent arbitrage* and some weaker form of *model-dependent arbitrage*. In particular, Davis and Hobson find that the expected dichotomy between the existence of a suitable martingale measure and the existence of a model-independent arbitrage does not hold in this specific setting; there can exist a third possibility in which there exists no suitable martingale measure but only model-dependent arbitrage opportunities (cf. [DH07, Def. 2.3]) can be constructed. A related notion of *weak arbitrage* is considered by Cox and Obłój [CO11b], where also the notion of *weak free lunch with vanishing risk (WFLVR)* [CO11b, Def. 2.1] is introduced in order to tackle the case of infinitely many given options. In the present paper we consider general path-dependent, possibly infinitely many, options and rule out the possibility of *weak* or *model-dependent arbitrage* by assuming that at least one option with super-linearly growing payoff is traded in the market. This is the key ingredient to obtain the model-free version of the *Fundamental Theorem of Asset Pricing* given in Theorem 1.3 which answers question (Q1).

In defining *arbitrage* we follow [DH07], where the concept of *model-independent arbitrage* is introduced in a very natural way, namely via semi-static strategies. A semi-static strategy consists of a static portfolio in finitely many options whose prices are known at time zero, and a dynamic, self-financing strategy in the underlying S . We say that a *model-independent arbitrage* exists when there is a semi-static portfolio

[†] University of Perugia, Department of Economics, Finance and Statistics, Via A. Pascoli 20, I-06123 Perugia

[‡] University of Vienna, Faculty of Mathematics, Nordbergstraße 15, A-1090 Wien.

with zero initial value and with strictly positive value at the terminal date. Strict positivity here pertains to *all* possible scenarios; there is no a priori reference measure to define a notion of *almost all* scenarios. Pioneering work in this regard was done by Hobson in [Hob98]; we refer to [Hob11, Section 2.6] for a detailed account of semi-static strategies and robust hedging. Cousot [Cou04, Cou07], Buehler [Bue06] and Carr and Madan [CM05] consider as given the prices of European call options and give, in different settings, necessary and sufficient conditions for the existence of calibrated arbitrage-free models. Davis, Obłój and Raval [DOR12, Theorem 3.6] tackle the case where a finite number of put options plus one additional European option with convex payoff is given; also the relevance for robust super-replication is discussed. In a one-period setting and assuming the prices of finitely many options, Riedel [Rie11] proves a robust Fundamental Theorem of Asset Pricing w.r.t. a weak notion of arbitrage.¹ In continuous time the situation is more delicate; for a discussion in this setting we refer to Cox and Obłój [CO11b] and Davis, Obłój and Raval [DOR12].

Heading for a Fundamental Theorem of Asset Pricing, the second issue concerns the pricing measures under consideration. Since we do not assume as given a reference measure, the obvious approach consists in considering as *admissible martingale measures* all probability measures on the path-space \mathbb{R}_+^T which are consistent with the observed option prices and under which the coordinate process is a martingale in its own filtration. In this setup we obtain Theorem 1.3, which connects the absence of arbitrage with the existence of an admissible pricing measure.

Having discussed this relation, it is natural to address the problem of super-replicating any other option written on S . The strategies used for replication again are of the semi-static kind described above. A central question is whether a model-free *Super-Replication Theorem* holds true: given a path-dependent derivative Φ , does the minimal endowment $p^R(\Phi)$ required for *super-replication* equal the *upper martingale price* $p^M(\Phi)$ obtained as the supremum of the expected value over admissible martingale measures? In a series of impressive achievements, Brown, Cox, Davis, Hobson, Klimmek, Madan, Neuberger, Obłój, Pederson, Raval, Rogers, Wang, Yor, and others [Rog93, Hob98, BHR01, HP02, MY02, CHO08, DOR12, CO11b, CO11a, CW12, HN12, HK12] were able to determine the values $p^R(\Phi)$ and $p^M(\Phi)$ explicitly for specific choices of Φ , showing in particular that they coincide. For an overview of the recent achievements we recommend the survey by Hobson [Hob11]. In the approach used by these authors, dominating tools are various Skorokhod-embedding techniques; we refer to the extensive overview given by Obłój in [Obł04]. In a discrete time setup, without assuming market-information, Deparis and Martini [DM04] establish the above duality for Φ satisfying a particular growth condition.²

Recently the super-replication problem in the model-free setting has been addressed via a new connection to the theory of optimal transport; see [GHLT11, TT11, BHL12]. In [GHLT11] Galichon, Henry-Labordère and Touzi systematically use a controlled stochastic dynamics approach, building on results of Tan and Touzi [TT11]. This enables the authors to derive the equation $p^M(\Phi) = p^R(\Phi)$ in the context of the look-back option when the terminal marginal of the underlying is known, recovering in particular results from [Hob98]. This viewpoint is developed further in [HLOST12] to include market information at intermediate times. In a discrete time setup the duality theory of optimal transport can be used to prove $p^R(\Phi) = p^M(\Phi)$ for general path-dependent Φ assuming knowledge on the intermediate marginals, see [BHL12]. In continuous time (assuming information on the terminal marginal) Dolinsky and Soner [DS12] are able to establish the relation $p^R(\Phi) = p^M(\Phi)$ for a large class of path-dependent derivatives. A robust super-replication result in a discrete time setting which also takes proportional transaction costs into account is established in [DS13].

In the present article, although inspired by the theory, we do not explicitly use results from optimal mass transport. Instead, we approach the Super-Replication Theorem using the classical route, i.e. through the Fundamental Theorem of Asset Pricing (Theorem 1.3). We obtain the relation $p^R(\Phi) = p^M(\Phi)$ under fairly

¹Under the assumption of a compact state-space, a Super-Replication Theorem is obtained as a corollary in [Rie11].

²We thank Marcel Nutz for bringing this paper to our attention.

general assumptions on the given market-information. In particular we recover the main result of [BHL12] as a special case.

Fundamental Theorem of Asset Pricing. We consider a finite, discrete time setting, with time horizon $T \in \mathbb{N}$, and a risky asset $S = (S_t)_{t=0}^T$, where S_0 is a positive real number which denotes the price of S to date. Formally, we take S to be the canonical process $S_t(x_1, \dots, x_T) = x_t$ on the path-space $\Omega = \mathbb{R}_+^T = [0, \infty)^T$.³ We also assume that there exists a risk free asset $B = (B_t)_{t=0}^T$ which is normalized to $B_t \equiv 1$. This setup allows for all possible choices of models since every non-negative stochastic process $S = (S_t)_{t=0}^T$ can be realized using the corresponding measure on the path-space.

Let I be some index set and $\varphi_i : \mathbb{R}_+^T \rightarrow \mathbb{R}$, $i \in I$, the payoff functions of options on the underlying S that can be *bought* on the market at time $t = 0$. W.l.o.g. we assume that they can be bought at price 0. We assume that, if an option φ can be both *bought and sold*, then bid and ask prices coincide. In this case we simply include $\pm\varphi$ among the φ_i . Consequently the set of admissible measures is defined as

$$\mathcal{P}_{(\varphi_i)_{i \in I}} := \left\{ \pi \in \mathcal{P}(\mathbb{R}_+^T) : \int_{\mathbb{R}_+^T} \varphi_i(x) d\pi(x) \leq 0, i \in I \right\}, \quad (1.1)$$

where $\mathcal{P}(\mathbb{R}_+^T)$ denotes the set of all probability measures on \mathbb{R}_+^T .

Definition 1.1 (Trading strategies). A trading strategy $\Delta = (\Delta_t)_{t=0}^{T-1}$ consists of Borel measurable functions $\Delta_t : \mathbb{R}_+^t \rightarrow \mathbb{R}$, where $0 \leq t < T$. The set of all such strategies will be denoted by \mathcal{H} . For the stochastic integral we use the notation

$$(\Delta \cdot x)_T := \sum_{t=0}^{T-1} \Delta_t(x_1, \dots, x_t)(x_{t+1} - x_t),$$

so that $(\Delta \cdot S)_T$ represents the gains or losses obtained by trading according to Δ .

The set of *martingale measures* \mathcal{M} consists of all probabilities on \mathbb{R}_+^T with finite first moment such that the canonical process S is a martingale in the natural filtration. Therefore, the set of admissible martingale measures is given by

$$\mathcal{M}_{(\varphi_i)_{i \in I}} := \mathcal{P}_{(\varphi_i)_{i \in I}} \cap \mathcal{M}. \quad (1.2)$$

As mentioned above, we define *arbitrage* via semi-static strategies, following [DH07, Def. 2.1].

Definition 1.2 (Arbitrage). There is model-independent arbitrage if there exists a trading strategy $\Delta \in \mathcal{H}$ and if there exist constants $a_1, \dots, a_N \geq 0$ and indices $i_1, \dots, i_N \in I$ such that

$$f(x_1, \dots, x_T) = \sum_{n=1}^N a_n \varphi_{i_n}(x_1, \dots, x_T) + (\Delta \cdot x)_T > 0 \quad (1.3)$$

for all $x_1, \dots, x_T \in \mathbb{R}_+$.

We emphasize the fact that *model-independent* arbitrage requires the strict inequality in (1.3) to hold *surely*, i.e., on the whole path-space \mathbb{R}_+^T .

In the Fundamental Theorem of Asset Pricing given below (Theorem 1.3) we assume the existence of an option with a super-linearly growing payoff $\varphi_0(S) = g(S_T)$ for some convex super-linear function $g : \mathbb{R}_+ \rightarrow \mathbb{R}$.

Theorem 1.3 (FTAP). Let $\varphi_i, i \in I$ be continuous functions on \mathbb{R}_+^T . Let $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a convex super-linear function and assume φ_0 to be of the form $\varphi_0(S) = g(S_T)$, where we suppose that 0 is an element of the index-set I . Assume also that

$$\lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^+}{m(x)} < \infty \quad \text{and} \quad \lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^-}{m(x)} = 0, \quad i \in I, \quad (1.4)$$

³We remark that the results obtained below are also valid in the case where S is allowed to take values on the whole real line. The proofs carry over to this setup without requiring significant changes.

where $m(x_1, \dots, x_T) := \sum_{t=1}^T g(x_t)$. Then the following are equivalent:

- (i) There is no model-independent arbitrage.
- (ii) $\mathcal{M}_{(\varphi_i)_{i \in I}} \neq \emptyset$.

Condition (1.4) is satisfied, for instance, when the set of the φ_i consists of European call options plus one power option.

Robust Super-Replication Results. As in classical mathematical finance, the Fundamental Theorem of Asset Pricing has a Super-Replication Theorem as immediate corollary.

Theorem 1.4 (Super-Replication). *Let $(\varphi_i)_{i \in I}$ be as in Theorem 1.3 and assume that $\mathcal{M}_{(\varphi_i)_{i \in I}} \neq \emptyset$. Let $\Phi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ be u.s.c. and such that*

$$\lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^+}{m(x)} = 0. \quad (1.5)$$

Then

$$p^M(\Phi) := \sup_{\pi \in \mathcal{M}_{(\varphi_i)_{i \in I}}} \int_{\mathbb{R}_+^T} \Phi(x) d\pi(x) \quad (1.6)$$

$$= \inf \left\{ d : \exists a_n \geq 0, \Delta \in \mathcal{H} \text{ s.t. } d + \sum_{n=1}^N a_n \varphi_{i_n} + (\Delta \cdot x)_T \geq \Phi \right\} =: p^R(\Phi). \quad (1.7)$$

In addition, the above supremum is a maximum.

We emphasize that the Super-Replication Theorem perfectly fits the setup of model-independent finance: the financial market provides information about the prices of traded derivatives φ_i , $i \in I$. This allows to access the largest reasonable price of the derivative Φ in two ways.

- (1) Following the no-arbitrage pricing paradigm, one selects (randomly or by another method) a martingale measure π which fits to the market prices; the corresponding price for the derivative Φ equals $\int_{\mathbb{R}_+^T} \Phi(x) d\pi(x)$. In general there are infinitely many possible choices for π and the robust point of view is to take (the) one leading to the largest value for $\int_{\mathbb{R}_+^T} \Phi(x) d\pi(x)$. This leads to $p^M(\Phi)$ given in (1.6).
- (2) On the other hand, a robust upper bound to the price of Φ can be obtained by considering semi-static super-hedges $d + \sum_{n=1}^N a_n \varphi_{i_n} + (\Delta \cdot x)_T \geq \Phi$. This approach was introduced by Hobson (cf. [Hob11]) and leads to the value $p^R(\Phi)$ in (1.7).

Theorem 1.4 asserts that the two approaches are equivalent.

The results presented so far required that the market sells a financial derivative $\varphi_0(S) = g(S_T)$ where g grows super-linearly. This assumption can be avoided, provided that a sufficient amount of call options written on S_T is traded on the market. For instance, it suffices to assume that there is a sequence of strikes K_n , $n \geq 1$, $K_n \rightarrow \infty$ such that the call options $\psi_{K_n} = (S_T - K_n)_+$ can be bought in the market at price p_n , where $p_n \rightarrow 0$ as $n \rightarrow \infty$. This is spelled out in detail in Corollary 4.2 below; in this introductory section we just present a particular consequence.

A prevalent assumption in the theory of model-independent pricing is that the distribution of S_T can be deduced from market data. This is due to the important observation of Breeden and Litzenberger [BL78] that knowing the *marginal* $S_T \sim \nu$ is essentially equivalent to knowing the prices p_K of $(S_T - K)_+$ for all strikes $K \geq 0$. The reasonable price of an arbitrary European derivative $\varphi(S_T)$ is then given by $\mathbb{E}_\nu[\varphi(S_T)] = \int_{\mathbb{R}_+} \varphi(y) d\nu(y)$. We write $\mathcal{M}(\nu)$ for the set of all martingale measures π satisfying $S_T(\pi) = \nu$. Of course this set is non-empty if and only if the first moment of ν exists and equals S_0 .

Corollary 1.5 (Super-Replication). *Assume that ν is a probability measure on \mathbb{R}_+ with finite first moment and barycenter S_0 . Let $\Phi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ be u.s.c. and linearly bounded from above. Then*

$$\begin{aligned} p^M(\Phi) &:= \sup_{\pi \in \mathcal{M}(\nu)} \left\{ \int_{\mathbb{R}_+^T} \Phi(x) d\pi(x) \right\} \\ &= \inf \left\{ \int_{\mathbb{R}_+} \varphi(y) d\nu(y) : \varphi \in L^1(\nu), \exists \Delta \in \mathcal{H} \text{ s.t. } \varphi(x_T) + (\Delta \cdot x)_T \geq \Phi(x) \right\} =: p^R(\Phi). \end{aligned}$$

In addition, the above supremum is a maximum.

More generally these results hold true if there exists a convex super-linear function $\tilde{g} : \mathbb{R}_+ \rightarrow \mathbb{R}$ in $L^1(\nu)$ such that

$$\lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^+}{\sum_{t=1}^T \tilde{g}(x_t)} < \infty. \quad (1.8)$$

In the same spirit we also recover [BHL12, Theorem 1] which corresponds to the Super-Replication Theorem in the particular case where all marginals $S_t \sim \mu_t$, $t = 1, \dots, T$ are known (see Corollary 4.5).

Concerning the just given super-replication results, a natural question is whether the infimum over super-replication strategies is in fact a minimum. This is in general not the case. In [BHL12, Section 4.3] a counterexample is constructed in a setup where $T = 2$ and the function Φ is uniformly bounded. As a remedy it may be useful to consider a *relaxed* notion of super-replication strategies. E.g., such “weak minimizers” are the critical tool in [BJ12, Appendix A]. We also refer to the recent article [Nut13] of Nutz; there the main focus is to obtain optimal super-replication strategies. The main difference to the present setup is that super-replication is understood w.r.t. a family of probability measures rather than in a path-wise sense.

Connection with Martingale Inequalities. Assume that Φ, φ are functions satisfying some proper integrability assumption. A path-wise hedging inequality of the form

$$\Phi(x_1, \dots, x_T) \leq \varphi(x_1, \dots, x_T) + (\Delta \cdot x)_T, \quad x_1, \dots, x_T \in \mathbb{R}_+, \quad (1.9)$$

implies that for every martingale $S = (S_t)_{t=1}^T$ we have

$$\mathbb{E}[\Phi(S_1, \dots, S_T)] \leq \mathbb{E}[\varphi(S_1, \dots, S_T)].$$

This follows by applying the inequality (1.9) to the paths of S and taking expectations. In short, every path-wise hedging inequality yields a martingale inequality as a direct consequence.

Conversely one may ask if a given martingale inequality can be established in this way, i.e. as a consequence of a path-wise hedging inequality of the form (1.9). In Section 5 below we explain why this can be expected as a consequence of the Super-Replication Theorem 1.4. Indeed an early version of this result motivated the path-wise approach to the Doob L^p -inequalities given in [ABP⁺12].

Organization of the paper. In Sections 2 and 3 we prove the Fundamental Theorem of Asset Pricing (Theorem 1.3) and the Super-Replication Theorem (Theorem 1.4), respectively. Section 4 collects different super-replication results which do not require the existence of super-linearly growing derivatives in the market. Finally we discuss the relation between robust Super-replication and Martingale Inequalities in Section 5.

2. FUNDAMENTAL THEOREM OF ASSET PRICING

In the definition of model-independent arbitrage we make use of trading strategies $\Delta \in \mathcal{H}$ which depend on S measurably but need not be bounded. In particular, $(\Delta \cdot S)$ is not necessarily integrable w.r.t. a martingale measure $\pi \in \mathcal{M}$. The following lemma allows us to take care of the thus arising subtleties.

Lemma 2.1. *Let $\pi \in \mathcal{M}$ and $\Delta \in \mathcal{H}$. Let $\varphi, \psi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ be measurable and such that $\int \varphi d\pi, \int \psi d\pi$ are well-defined⁴. Then*

$$\varphi(x) + (\Delta \cdot x)_T \geq \psi(x), \quad x \in \mathbb{R}_+^T \quad \implies \quad \int \varphi d\pi \geq \int \psi d\pi. \quad (2.1)$$

If at least one of the integrals $\int \varphi d\pi, \int \psi d\pi$ is finitely valued then we also have

$$\varphi(x) + (\Delta \cdot x)_T > \psi(x), \quad x \in \mathbb{R}_+^T \quad \implies \quad \int \varphi d\pi > \int \psi d\pi. \quad (2.2)$$

Proof. To prove (2.1) we may assume that $\int \varphi^+ d\pi < \infty, \int \psi^- d\pi < \infty$, the other cases are trivial. It follows that also $\int \varphi - \psi d\pi$ exists. Hence by replacing φ with $\varphi - \psi$ we may assume that $\psi \equiv 0$. Assume that $T = 2$, set $\mu := S_1(\pi)$ and consider the disintegration $(\pi_{x_1})_{x_1 \in \mathbb{R}_+}$ of π with respect to μ . Then we have

$$\int \varphi d\pi = \int \left(\int \varphi(x_1, x_2) d\pi_{x_1}(x_2) \right) \quad (2.3)$$

$$= \int \left(\int \varphi(x_1, x_2) + \Delta(x_1)(x_2 - x_1) d\pi_{x_1}(x_2) \right) \geq 0. \quad (2.4)$$

In the general case $T \in \mathbb{N}$ one iterates this argument $T - 1$ times; we omit this.

The proof of (2.2) works in the same way. □

As a particular consequence of Lemma 2.1 the existence of a martingale measure in $\mathcal{M}_{(\varphi_i)_{i \in I}}$ implies that there is no model-independent arbitrage.

Proof of Theorem 1.3, (ii) \implies (i). Pick $\pi \in \mathcal{M}_{(\varphi_i)_{i \in I}}$ and assume that there exists $f(x) = \sum_{n=1}^N a_n \varphi_{i_n}(x) + (\Delta \cdot x)_T$, where $a_n \geq 0$ and $\Delta \in \mathcal{H}$ such that $f > 0$. Applying (2.2) to $\varphi = \sum_{n=1}^N a_n \varphi_{i_n}(x)$ and $\psi \equiv 0$ then gives the desired contradiction. □

In the same fashion Lemma 2.1 yields the “intuitively obvious” inequality $p^M(\Phi) \leq p^R(\Phi)$ in our super-replication results.

It is of course natural to ask why we do not only consider bounded strategies. We explain here why this would be too restrictive for our purposes. For every convex function $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ and $x_t, x_{t+1} \in \mathbb{R}_+$ we have⁵

$$g(x_t) + g'(x_t)(x_{t+1} - x_t) \leq g(x_{t+1}). \quad (2.5)$$

This simple inequality expresses a fact which is widely known in finance: a convex derivative written on S_t can be super-replicated using the corresponding derivative written on S_{t+1} . To incorporate this argument in our path-wise hedging framework, we need to include $\Delta_t(x_1, \dots, x_t) := g'(x_t)$ in the set of admissible trading strategies.

Indeed, in showing the non trivial implication (i) \implies (ii) in of Theorem 1.3 (and the non-trivial inequality $p^M(\Phi) \geq p^R(\Phi)$ in our Super-Replication Theorems), it is sufficient to use the no arbitrage assumption on a subset of \mathcal{H} which consists entirely of strategies Δ such that $(\Delta \cdot S)$ is π -integrable for all $\pi \in \mathcal{M}_{(\varphi_i)_{i \in I}}$.

Definition 2.2 (*g-Admissible Strategy*). *A trading strategy $\Delta = (\Delta_t)_{t=0}^{T-1}$ is called g-admissible if, for $0 \leq t \leq T - 1$, $\Delta_t : \mathbb{R}_+^t \rightarrow \mathbb{R}$ is a continuous function such that, for some $c \in \mathbb{R}_+$,*

$$|\Delta_t(x_1, \dots, x_t)(x_{t+1} - x_t)| \leq c \left(1 \vee \sum_{s=1}^{t+1} g(x_s) \right). \quad (2.6)$$

The set of all g-admissible trading strategies is denoted by \mathcal{H}_g .

⁴We say that $\int f d\pi$ is well-defined if $\int f^+ d\pi < \infty$ or $\int f^- d\pi < \infty$.

⁵In the (at most countably many) points where the convex function g is not differentiable, we define g' as its right derivative.

Trivially we have $\mathcal{H}_g \subseteq \mathcal{H}$. We briefly comment on the integrability properties of the set \mathcal{H}_g . Assume that π is a martingale measure on \mathbb{R}_+^T such that $\int g(x_T) d\pi(x) < \infty$. By Jensen's inequality we then have $\int g(x_t) d\pi(x) < \infty$ also for all $t < T$. Thus for $\Delta \in \mathcal{H}_g$, (2.6) implies that

$$\int_{\mathbb{R}_+^T} |\Delta_t(x_1, \dots, x_t)(x_{t+1} - x_t)| d\pi(x) < \infty.$$

Disintegrating π w.r.t. (x_1, \dots, x_t) it moreover follows that

$$\int_{\mathbb{R}_+^T} \Delta_t(x_1, \dots, x_t)(x_{t+1} - x_t) d\pi(x) = 0.$$

Note also that by (2.5) $|g'(x_t)(x_{t+1} - x_t)| \leq |g(x_t)| + |g(x_{t+1})|$, hence $\Delta_t(x_1, \dots, x_t) := g'(x_t)$ is g -admissible.

In the following proposition we use the notation introduced in (1.1) for the set of admissible measures. Recall that we write $m(x_1, \dots, x_T) = \sum_{t=1}^T g(x_t)$.

Proposition 2.3. *Let $\varphi_i : \mathbb{R}_+^T \rightarrow \mathbb{R}$, $i = 1, \dots, N$ be continuous functions satisfying*

$$\lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^+}{m(x)} < \infty \quad \text{and} \quad \lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^-}{m(x)} = 0 \quad (2.7)$$

and set $\varphi_{N+1} := m \vee 1 =: \bar{m}$. TFAE:

(i) *There is no $f = \sum_{n=1}^{N+1} a_n \varphi_n$ with $a_n \geq 0$ s.t.*

$$f(x) > 0 \quad \text{for all } x \in \mathbb{R}_+^T.$$

(i') *There is no $f = \sum_{n=1}^{N+1} a_n \varphi_n$ with $a_n \geq 0$ s.t.*

$$f(x) \geq \bar{m}(x) \quad \text{for all } x \in \mathbb{R}_+^T.$$

(ii) $\mathcal{P}_{(\varphi_i)_{i=1}^{N+1}} \neq \emptyset$.

Proof. The only non trivial implication is (i') \Rightarrow (ii): Consider the Banach space $C_m^b(\mathbb{R}_+^T)$ of continuous functions f on \mathbb{R}_+^T such that

$$\|f\|_{C_m^b} = \sup_{x \in \mathbb{R}_+^T} \frac{|f(x)|}{\bar{m}(x)} < \infty.$$

The norm is designed in such a way that the multiplication operator $T_m : C_m^b(\mathbb{R}_+^T) \rightarrow C^b(\mathbb{R}_+^T)$

$$T_m(f) = \frac{f}{\bar{m}}$$

is an isometry, where the Banach space $C^b(\mathbb{R}_+^T)$ of bounded continuous functions h on \mathbb{R}_+^T is endowed with

$$\|h\|_{C^b} = \sup_{x \in \mathbb{R}_+^T} |h(x)| < \infty.$$

Recall that $C^b(\mathbb{R}_+^T)$ may be identified with the space $C(\check{\mathbb{R}}_+^T)$ of continuous functions on the Stone-Cech-compactification $\check{\mathbb{R}}_+^T$ of \mathbb{R}_+^T . Hence the dual space of $C^b(\mathbb{R}_+^T)$ can be identified with $\mathcal{M}(\check{\mathbb{R}}_+^T)$, the space of signed Radon measures μ on $\check{\mathbb{R}}_+^T$. Each μ can be uniquely decomposed into $\mu = \mu^r + \mu^s$, where the regular part μ^r is supported by \mathbb{R}_+^T while the singular part μ^s is supported by $\check{\mathbb{R}}_+^T \setminus \mathbb{R}_+^T$. The bottom line of these considerations is that a continuous linear functional F on $(C_m^b(\mathbb{R}_+^T), \|\cdot\|_{C_m^b})$ is given by some $\mu = \mu^r + \mu^s \in \mathcal{M}(\check{\mathbb{R}}_+^T)$ via

$$\begin{aligned} F(f) &= \int \frac{f(x)}{\bar{m}(x)} d\mu(x) \\ &= \int \frac{f(x)}{\bar{m}(x)} d\mu^r(x) + \int \frac{f(x)}{\bar{m}(x)} d\mu^s(x), \quad \text{for } f \in C_m^b(\mathbb{R}_+^T). \end{aligned} \quad (2.8)$$

Finally observe that the interior of the positive orthant of $C_m^b(\mathbb{R}_+^T)$ is given by

$$(C_m^b)_{++}(\mathbb{R}_+^T) = \left\{ f \in C_m^b : \inf_{x \in \mathbb{R}_+^T} \frac{f(x)}{\bar{m}(x)} > 0 \right\},$$

as one easily sees from the isometric identification of $C_m^b(\mathbb{R}_+^T)$ with $C(\check{\mathbb{R}}_+^T)$.

Turning to the present setting, define K as the compact, convex set in $C_m^b(\mathbb{R}_+^T)$

$$K := \left\{ \sum_{n=1}^{N+1} a_n \varphi_n : a_n \geq 0, \sum_{n=1}^{N+1} a_n = 1 \right\}.$$

By assumption (i') we have

$$K \cap (C_m^b)_{++}(\mathbb{R}_+^T) = \emptyset,$$

so that we may apply Hahn-Banach to find a linear functional $F \in C_m^b(\mathbb{R}_+^T)^*$ separating K from $(C_m^b)_{++}(\mathbb{R}_+^T)$, i.e. some $\mu = \mu^r + \mu^s \in \mathcal{M}(\check{\mathbb{R}}_+^T)$ such that

$$\int \frac{f(x)}{\bar{m}(x)} d\mu(x) > 0 \quad \text{for all } f \in (C_m^b)_{++}(\mathbb{R}_+^T), \quad (2.9)$$

while

$$\int \frac{f(x)}{\bar{m}(x)} d\mu(x) \leq 0 \quad \text{for all } f \in K. \quad (2.10)$$

Clearly (2.9) implies that $\mu = \mu^r + \mu^s$ is positive. We first observe that we have $\mu^r \neq 0$. Indeed, supposing $\mu^r = 0$, we find

$$\int \frac{\varphi_{N+1}(x)}{\bar{m}(x)} d\mu(x) = \int \frac{\varphi_{N+1}(x)}{\bar{m}(x)} d\mu^s(x) = \int 1 d\mu^s(x) = \|\mu^s\| > 0$$

and this is in contradiction to (2.10).

We now claim that μ^r also separates K from $(C_m^b)_{++}(\mathbb{R}_+^T)$. On the one hand, μ^r is a positive measure on \mathbb{R}_+^T . Hence (2.9) still holds true, with μ replaced by μ^r . On the other hand, for each $1 \leq n \leq N+1$, we have

$$\int \frac{\varphi_n(x)}{\bar{m}(x)} d\mu^r(x) \leq \int \frac{\varphi_n(x)}{\bar{m}(x)} d\mu(x) \leq 0.$$

The second inequality follows from (2.10). For the first inequality it suffices to remark that

$$\int \frac{\varphi_n(x)^-}{\bar{m}(x)} d\mu^s(x) = 0, \quad n = 1, \dots, N+1,$$

by (2.7). By normalizing μ^r to $\pi := \frac{\mu^r}{\|\mu^r\|}$, we find a positive probability π on \mathbb{R}^T with $\int \frac{\varphi_n(x)}{\bar{m}(x)} d\pi(x) \leq 0$, for $n = 1, \dots, N+1$. Now define $\hat{\pi}$ by

$$\frac{d\hat{\pi}}{d\pi} = \frac{1}{\bar{m}} \left(\int \frac{1}{\bar{m}} d\pi \right)^{-1}.$$

We have that $\hat{\pi}$ is a positive probability on \mathbb{R}^T with $\int \varphi_n d\hat{\pi} \leq 0$, for $n = 1, \dots, N+1$, which shows that $\mathcal{P}_{(\varphi_i)_{i=1}^{N+1}} \neq \emptyset$. \square

The above proposition is the basis for the proof of the non-trivial part of Theorem 1.3. In the course of the argument we also use the following characterization of martingale measures.

$$\mathcal{M} = \left\{ \pi \in \mathcal{P}(\mathbb{R}_+^T) : \begin{array}{l} S_t \text{ has finite first moment w.r.t. } \pi, t \leq T \\ \int_{\mathbb{R}_+^T} (\Delta \cdot x)_T d\pi(x) = 0, \Delta \in C_b, \end{array} \right\}, \quad (2.11)$$

where with $\Delta \in C_b$ we mean that Δ_t is continuous and bounded for all $t = 0, \dots, T-1$. The proof of (2.11) is straightforward, see for instance [BHL12].

Proof of Theorem 1.3, (i) \Rightarrow (ii). In fact here we prove a stronger result, that is, we show that $(i)^* \Rightarrow (ii)$, where condition $(i)^*$ is defined as

$(i)^*$ There is no model-independent arbitrage such that $\Delta \in \mathcal{H}_g$ (see Definition 2.2).

Recall that $\varphi_0(x_1, \dots, x_T) = g(x_T)$ and set

$$\varphi_{-1}(x_1, \dots, x_T) := - \sum_{t=1}^{T-1} g'(x_t)(x_T - x_t) + Tg(x_T).$$

Note that since no arbitrage strategy can be constructed using the option φ_0 , and since $g'(x_t), t < T$ are g -admissible trading strategies, it follows that no arbitrage strategy can be constructed with the help of φ_{-1} . We make the crucial observation that due to the convexity of g we have $m \leq \varphi_{-1}$. Moreover, if π is a martingale measure, then $\int_{\mathbb{R}_+^T} \varphi_{-1} d\pi = \int_{\mathbb{R}_+^T} T\varphi_0 d\pi$. Note that $\mathcal{M}_{(\varphi_i)_{i \in I}} = \mathcal{M}_{(\varphi_i)_{i \in I}, m}$ by Jensen's inequality. We will use a compactness argument to show that this set is not empty.

Assume that we are given finite families F_1, F_2 , where $F_1 \subseteq I$ and

$$\{\varphi_i\}_{i \in F_2} \subseteq \{\Delta_t(x_1, \dots, x_t)(x_{t+1} - x_t) : t < T, \Delta_t \in C_b(\mathbb{R}_+^t)\}.$$

Then there exists no arbitrage, in the sense of Proposition 2.3, for the family

$$\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\} \cup \{-1\}}. \quad (2.12)$$

Since $m \leq \varphi_{-1}$ there is still no arbitrage opportunity if we replace φ_{-1} by m . Since the functions Δ_t in (2.12) are taken to be continuous and bounded we may apply Proposition 2.3 to the family

$$\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}$$

to obtain that

$$\mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m} \neq \emptyset.$$

Since

$$\mathcal{M}_{(\varphi_i)_{i \in I}} = \bigcap_{F_1, F_2} \mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m},$$

it remains to prove that $\mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$ is compact.

Step 1. Relative compactness.

We show that the set $\mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$ is tight, hence relatively compact by Prokhorov's theorem. First we recall that $\lim_{\|x\| \rightarrow \infty} \frac{m(x)}{\|x\|} = \infty$ and that $\int_{\mathbb{R}_+^T} m d\pi \leq 0$ for $\pi \in \mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$. This implies $-\infty < -a := \min m < 0$ and that for all δ there is k_δ s.t. $m > \frac{1}{\delta}$ on K_δ^c , where $K_\delta := [0, k_\delta]^T$. Hence

$$\int_{K_\delta^c} m d\pi \geq \frac{1}{\delta} \pi(K_\delta^c). \quad (2.13)$$

Furthermore

$$0 \geq \int_{\mathbb{R}_+^T} m d\pi = \int_{K_\delta} m d\pi + \int_{K_\delta^c} m d\pi \geq -a\pi(K_\delta) + \int_{K_\delta^c} m d\pi,$$

that is,

$$\int_{K_\delta^c} m d\pi \leq a\pi(K_\delta). \quad (2.14)$$

Putting things together we obtain

$$\pi(K_\delta^c) \leq \delta \int_{K_\delta^c} m d\pi \leq \delta a\pi(K_\delta) \leq \delta a.$$

This proves that for each fixed $\varepsilon > 0$ there is $k (= k_\delta$ for $\delta = \varepsilon/a)$ such that $\pi([0, k]^T)^c \leq \varepsilon$ for all $\pi \in \mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$. Hence, $\mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$ is tight and thus relatively compact by Prokhorov's theorem.

Step 2. Closedness.

Let $\pi_n \in \mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$ be such that (π_n) converges weakly to $\tilde{\pi}$. We are going to prove that $\tilde{\pi} \in \mathcal{P}_{\{\varphi_i\}_{i \in F_1 \cup F_2 \cup \{0\}}, m}$. Since $\tilde{\pi}$ is clearly a probability measure, we only need to prove that $\tilde{\pi}$ satisfies the admissibility constraints:

$$\int_{\mathbb{R}_+^T} \varphi \, d\tilde{\pi} \leq 0, \quad \varphi \in \{m, \varphi_i : i \in F_1 \cup F_2 \cup \{0\}\}.$$

We will consider separately the two integrals $\int_{\mathbb{R}_+^T} \varphi^+ \, d\tilde{\pi}$ and $\int_{\mathbb{R}_+^T} \varphi^- \, d\tilde{\pi}$.

First of all, for each $\varphi \in \{m, \varphi_i : i \in F_1 \cup F_2 \cup \{0\}\}$ and for every $u \in [0, \infty)$ we have the basic inequality

$$\int_{\mathbb{R}_+^T} \varphi^+ \, d\pi_n \geq \int_{\mathbb{R}_+^T} \varphi^+ \wedge u \, d\pi_n.$$

Taking the limit $n \rightarrow \infty$ and then the limit $u \rightarrow \infty$ on both sides, we obtain

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^+ \, d\pi_n \geq \lim_{u \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^+ \wedge u \, d\pi_n = \lim_{u \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^+ \wedge u \, d\tilde{\pi} = \int_{\mathbb{R}_+^T} \varphi^+ \, d\tilde{\pi}, \quad (2.15)$$

by weak convergence and by monotone convergence.

Furthermore, we will show that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^- \, d\pi_n = \int_{\mathbb{R}_+^T} \varphi^- \, d\tilde{\pi}. \quad (2.16)$$

Inequality (2.15) and equation (2.16) together then yield

$$\int_{\mathbb{R}_+^T} \varphi \, d\tilde{\pi} = \int_{\mathbb{R}_+^T} \varphi^+ \, d\tilde{\pi} - \int_{\mathbb{R}_+^T} \varphi^- \, d\tilde{\pi} \leq \lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^+ \, d\pi_n - \lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^- \, d\pi_n = \lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi \, d\pi_n \leq 0,$$

as wanted.

In order to prove (2.16) we will use the previous step, that is, for any fixed $\varepsilon > 0$ there is $k = k_\varepsilon > 0$ such that $\pi_n(K^c) \leq \varepsilon$ for all $n \in \mathbb{N}$, where $K := [0, k]^T$. By definition of weak convergence we have

$$\lim_{n \rightarrow \infty} \int_K \varphi^- \, d\pi_n = \int_K \varphi^- \, d\pi. \quad (2.17)$$

Therefore, if $(k_\varepsilon)_\varepsilon$ is bounded, then we are done. We hence suppose that $k_\varepsilon \rightarrow \infty$ as $\varepsilon \rightarrow 0$. Note that $\int_{\mathbb{R}_+^T} m \, d\pi_n \leq 0$ gives $\int_{\mathbb{R}_+^T} (m + a + 1) \, d\pi_n \leq a + 1$, which in turn implies $\int_A (m + a + 1) \, d\pi_n \leq a + 1$ for every $A \subseteq \mathbb{R}_+^T$, being $m + a + 1$ non-negative (actually, $m + a + 1 \geq 1$). Thus for $\varphi \in \{m, \varphi_i : i \in F_1 \cup F_2 \cup \{0\}\}$ we have

$$a + 1 \geq \int_{K^c} (m + a + 1) \mathbb{1}_{\varphi^- > 0} \, d\pi_n \geq \int_{K^c} \varphi^- \min_{K^c} \frac{(m + a + 1)}{\varphi^-} \mathbb{1}_{\varphi^- > 0} \, d\pi_n,$$

which implies

$$\int_{K^c} \varphi^- \, d\pi_n = \int_{K^c} \varphi^- \mathbb{1}_{\varphi^- > 0} \, d\pi_n \leq (a + 1) \max_{K^c} \frac{\varphi^-}{(m + a + 1)}.$$

Now note that for all $\varphi \in \{m, \varphi_i : i \in F_1 \cup F_2 \cup \{0\}\}$ we have that $\max_{K^c} \frac{\varphi^-}{(m + a + 1)} \rightarrow 0$ as $\varepsilon \rightarrow 0$. From this it follows that

$$\lim_{n \rightarrow \infty} \int_{K^c} \varphi^- \, d\pi_n = 0 \quad (2.18)$$

as $\varepsilon \rightarrow 0$, uniformly in n . Together, (2.17) and (2.18) imply that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^T} \varphi^- \, d\pi_n = \int_{\mathbb{R}_+^T} \varphi^- \, d\tilde{\pi},$$

as claimed. This concludes the proof. \square

Remark 2.4. *If the stock prices process is not allowed to take values on the (whole) half-line \mathbb{R}_+ but is restricted to a bounded interval $[0, b]$ then the above considerations simplify significantly. In this case the*

path-space is compact, all continuous functions φ_i are bounded and the set of admissible measures is automatically compact; there is no need to require the existence of options whose payoff grows super-linearly. As a consequence, the robust FTAP follows in a straightforward way from the Hahn-Banach Theorem.

3. SUPER-REPLICATION THEOREM

The Super-Replication Theorem 1.4 is a direct consequence of the Fundamental Theorem of Asset Pricing, Theorem 1.3.

Proof of Theorem 1.4. By Lemma 2.1 $p^M(\Phi) \leq p^R(\Phi)$. It remains to prove the converse inequality. In fact we prove a result which is stronger than the one stated. That is, we show this inequality when using only g -admissible strategies in the dual problem, i.e., when replacing \mathcal{H} by \mathcal{H}_g in the minimization problem in (1.7). Let us first consider the case of continuous Φ satisfying (1.5) and

$$\lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^-}{m(x)} < \infty. \quad (3.1)$$

Now suppose that the inequality is strict, that is, there exists p such that

$$p^M(\Phi) < p < p^R(\Phi). \quad (3.2)$$

Define $\varphi := -\Phi + p$ and note that Theorem 1.3 applies to the set of constraints $\{\varphi, \varphi_i, i \in I\}$, implying the equivalence of the following:

- (i) $\nexists f(x) = \sum_{n=1}^N a_n \varphi_{i_n}(x) + \varphi(x) + (\Delta \cdot x)_T > 0$ with $a_n \geq 0$ and $\Delta \in \mathcal{H}_g$,
- (ii) $\mathcal{M}_{(\varphi_i)_{i \in I}, \varphi} \neq \emptyset$.

Therefore, either there exists $\pi \in \mathcal{M}_{(\varphi_i)_{i \in I}}$ such that

$$\int_{\mathbb{R}_+^T} \Phi d\pi \geq p, \quad (3.3)$$

or there exist $a_n \geq 0$ and $\Delta \in \mathcal{H}_g$ such that

$$p + \sum_{n=1}^N a_n \varphi_{i_n}(x) + (\Delta \cdot x)_T > \Phi(x). \quad (3.4)$$

Note that (3.3) would imply $p^M(\Phi) \geq p$, in contradiction to the first inequality in (3.2), and that (3.4) would imply $p^R(\Phi) \leq p$, in contradiction to the second inequality in (3.2). This shows that there is no p as in (3.2), hence the duality stated in the theorem holds for all continuous Φ which satisfy (1.5) and (3.1). Now note that any u.s.c. function Φ satisfying (1.5) can be written as an infimum over continuous functions $\Phi_n, n \in \mathbb{N}$ satisfying (1.5) and (3.1). By a standard argument, the duality relation then carries over from Φ_n to Φ . This is worked out in detail for instance in [BHL12, Theorem 1] in a very similar setup. At the same place the reader can find the argument showing that the supremum in (1.6) is attained. \square

4. RAMIFICATIONS OF THE SUPER-REPLICATION RESULT

We start with a corollary of the previous results which avoids the asymmetry present in the requirements on φ_0 . To achieve this, we assume that there exists a sequence of call options written on S_T whose strikes K_n tend to ∞ . We call p_n the corresponding market prices and use the notation

$$\psi_n(y) := (y - K_n)_+, \quad \tilde{\psi}_n(x) := (\psi_n(x_T) - p_n), \quad n \geq 1. \quad (4.1)$$

Assumption 4.1. Let $\varphi_i : \mathbb{R}_+^T \rightarrow \mathbb{R}, i \in I$ be continuous functions including $\tilde{\psi}_n, n \geq 1$, and assume that $\mathcal{M}_{(\varphi_i)_{i \in I}} \neq \emptyset$. Let $\Phi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ be u.s.c. and $\alpha_n \geq 0$ be such that $\sum_{n=1}^{\infty} \alpha_n = \infty$ and $\sum_{n=1}^{\infty} \alpha_n p_n < \infty$. We set

$$g_0(y) := \sum_{n=1}^{\infty} \alpha_n (\psi_n(y) - p_n), \quad m_0(x) := \sum_{t=1}^T g_0(x_t)$$

and assume that

$$\lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^-}{m_0(x)} = 0, \quad \lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^+}{m_0(x)} < \infty, \quad \lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^+}{m_0(x)} = 0.$$

Theorem 1.4 can then be applied by setting $g = g_0$. Indeed, since $\tilde{\psi}_n, n \geq 1$ are already present in the admissibility resp. the super-replication condition, it makes no difference whether or not one includes also g_0 among the φ_i . Hence we obtain:

Corollary 4.2. *Let $(\varphi_i)_{i \in I}$, Φ satisfy Assumption 4.1. Then*

$$\begin{aligned} p^M(\Phi) &:= \sup_{\pi \in \mathcal{M}_{(\varphi_i)_{i \in I}}} \int_{\mathbb{R}_+^I} \Phi(x) d\pi(x) \\ &= \inf \left\{ d : \exists a_1, \dots, a_N \geq 0, i_1, \dots, i_N \in I, b_n \geq 0, \sup_n b_n \alpha_n < \infty, \Delta \in \mathcal{H} \text{ s.t.} \right. \\ &\quad \left. d + \sum_{k=1}^N a_k \varphi_{i_k} + \sum_{n=1}^{\infty} b_n \alpha_n \tilde{\psi}_n + (\Delta \cdot x)_T \geq \Phi \right\} =: p^R(\Phi). \end{aligned}$$

In addition, the above supremum is a maximum.

Note that this result can be easily put in “symmetric form” including also $-\tilde{\psi}_n, n \geq 1$ in the family $(\varphi_i)_{i \in I}$. Moreover, as a consequence of Corollary 4.2, we have the super-replication result under the assumption that the distribution ν of the asset at the terminal date T is known. Here I is allowed to be the empty set, in which case we obtain exactly Corollary 1.5.

We note that, for any convex super-linear function $\bar{g} : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that $\int_{\mathbb{R}_+} \bar{g} d\nu < \infty$, there exist constants $c, \alpha_n \geq 0$ and $K_n \nearrow \infty$ such that

$$\bar{g}(y) \leq c + \sum_{n=1}^{\infty} \alpha_n (y - K_n)_+, \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} \alpha_n p_n < \infty, \quad (4.2)$$

where $p_n := \int_{K_n}^{\infty} (y - K_n) d\nu(y)$. Setting

$$\mathcal{M}_{(\varphi_i)_{i \in I}}(\nu) := \mathcal{M}_{(\varphi_i)_{i \in I}} \cap \mathcal{M}(\nu),$$

we obtain the following result.

Corollary 4.3. *Let $\varphi_i : \mathbb{R}_+^T \rightarrow \mathbb{R}, i \in I$ be continuous and growing at most linearly at infinity and assume $\mathcal{M}_{(\varphi_i)_{i \in I}}(\nu) \neq \emptyset$. For $\Phi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ u.s.c. and linearly bounded from above we have*

$$\begin{aligned} p^M(\Phi) &:= \sup_{\mathcal{M}_{(\varphi_i)_{i \in I}}(\nu)} \left\{ \int_{\mathbb{R}_+^T} \Phi(x) d\pi(x) \right\} \\ &= \inf \left\{ \int_{\mathbb{R}_+} \varphi(y) d\nu(y) : \begin{array}{l} \varphi \in L^1(\nu), \exists \Delta \in \mathcal{H}, a_1, \dots, a_N \geq 0, i_1, \dots, i_N \in I, \\ \text{s.t. } \varphi(x_T) + \sum_{n=1}^N a_n \varphi_{i_n}(x) + (\Delta \cdot x)_T \geq \Phi(x) \end{array} \right\} =: p^R(\Phi). \end{aligned} \quad (4.3)$$

In addition, the above supremum is a maximum.

More generally these results hold true for $\varphi_i, i \in I$ continuous and Φ u.s.c. if there exists a convex super-linear function $\tilde{g} : \mathbb{R}_+ \rightarrow \mathbb{R}$ in $L^1(\nu)$ such that

$$\lim_{\|x\| \rightarrow \infty} \frac{|\varphi_i(x)|}{\sum_{t=1}^T \tilde{g}(x_t)} < \infty, \quad \lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^+}{\sum_{t=1}^T \tilde{g}(x_t)} < \infty. \quad (4.4)$$

Proof. Step 1. Let $\varphi_i, i \in I$ be continuous and such that $\mathcal{M}_{(\varphi_i)_{i \in I}}(\nu) \neq \emptyset$, and Φ be u.s.c. and such that (4.4) holds for some convex super-linear function $\tilde{g} \in L^1(\nu)$. By applying Lemma 4.4 to $f = \tilde{g}$, we obtain a convex super-linear function \bar{g} in $L^1(\nu)$ such that

$$\lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^-}{\sum_{t=1}^T \bar{g}(x_t)} = 0, \quad \lim_{\|x\| \rightarrow \infty} \frac{\varphi_i(x)^+}{\sum_{t=1}^T \bar{g}(x_t)} = 0, \quad \lim_{\|x\| \rightarrow \infty} \frac{\Phi(x)^+}{\sum_{t=1}^T \bar{g}(x_t)} = 0.$$

Now consider $\alpha_n \geq 0$ and $K_n \nearrow \infty$ as in (4.2). We can include the corresponding functions $\tilde{\psi}_n$ defined as in (4.1) among the φ_i since this neither changes the set of admissible martingale measures nor introduces

arbitrage. Now, applying Corollary 4.2 we obtain

$$\begin{aligned} \sup_{\mathcal{M}_{(\varphi_i)_{i \in I}}(\nu)} \left\{ \int_{\mathbb{R}_+^T} \Phi(x) d\pi(x) \right\} &= \inf \left\{ d : \begin{array}{l} \exists \Delta \in \mathcal{H}, a_1, \dots, a_N \geq 0, i_1, \dots, i_N \in I, b_n \geq 0, \sup_n b_n \alpha_n < \infty \\ \text{s.t. } d + \sum_{n=1}^N a_n \varphi_{i_n}(x) + \sum_{n=1}^{\infty} b_n \alpha_n \tilde{\psi}_n + (\Delta \cdot x)_T \geq \Phi(x) \end{array} \right\} \\ &\geq \inf \left\{ \int_{\mathbb{R}_+} \varphi(y) d\nu(y) : \begin{array}{l} \varphi \in L^1(\nu), \exists \Delta \in \mathcal{H}, a_1, \dots, a_N \geq 0, i_1, \dots, i_N \in I \\ \text{s.t. } \varphi(x_T) + \sum_{n=1}^N a_n \varphi_{i_n}(x) + (\Delta \cdot x)_T \geq \Phi(x) \end{array} \right\}. \end{aligned}$$

This gives $p^M(\Phi) \geq p^R(\Phi)$, hence $p^M(\Phi) = p^R(\Phi)$, the other inequality being trivial. The fact that the supremum in (4.3) is attained is again obtained by standard arguments, cf. [BHL12, Theorem 1].

Step 2. Let now $\varphi_i, i \in I$ be continuous and growing at most linearly at infinity and Φ be u.s.c. and linearly bounded from above. By applying Lemma 4.4 to $f(x) = |x|$, we obtain a convex super-linear function \tilde{f} in $L^1(\nu)$ such that (4.4) is satisfied with $\tilde{g} = \tilde{f}$. Now we can apply Step 1, which concludes the proof. \square

Lemma 4.4. *Let μ be a probability measure on \mathbb{R}_+ and $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ a convex function in $L^1(\mu)$. Then there exists a convex function $\tilde{f} : \mathbb{R}_+ \rightarrow \mathbb{R}$ in $L^1(\mu)$ such that $\frac{|\tilde{f}(x)|}{|f(x)|} \rightarrow \infty$ as $x \rightarrow \infty$.*

Proof. W.l.o.g. let $\int_{\mathbb{R}_+} f(x) d\mu(x) = 1$. Define

$$a_n := \int_{\mathbb{R}_+} \mathbb{1}_{f(x) \in [n, n+1)} f(x) d\mu(x),$$

so that $\sum_{n \in \mathbb{N}} a_n = 1$. For $k \in \mathbb{N}$, denote by N_k the set of all $n \in \mathbb{N}$ such that

$$\sum_{m \geq n} a_m \in \left[\frac{1}{2^{k+1}}, \frac{1}{2^k} \right)$$

and set $b_n := k + 1$ for all $n \in N_k$. This gives

$$\sum_{n \in \mathbb{N}} a_n b_n = \sum_{k \in \mathbb{N}} \sum_{n \in N_k} (k+1) a_n \leq \sum_{k \in \mathbb{N}} (k+1) \frac{1}{2^k} < \infty.$$

Now define the convex function \tilde{f} by $\tilde{f}(x) = \sum_{n \in \mathbb{N}} \mathbb{1}_{f(x) \in [n, n+1)}(x) f(x) a_n$, so that

$$\frac{|\tilde{f}(x)|}{|f(x)|} = \sum_{n \in \mathbb{N}} \mathbb{1}_{f(x) \in [n, n+1)}(x) a_n \rightarrow \infty \quad \text{for } x \rightarrow \infty$$

and $\int_{\mathbb{R}_+} \tilde{f}(x) d\mu(x) = \sum_{n \in \mathbb{N}} a_n b_n < \infty$. \square

It seems natural to assume that the market does not only yield information about the call options at the terminal time T . In fact, in [BHL12] a super-replication result is proved for the case where all marginals $S_t \sim \nu_t, t = 1, \dots, T$ are known. By Theorem 1 in [BHL12] we have:

Corollary 4.5. *Assume that $\nu_t, t = 1, \dots, T$ are probability measures on \mathbb{R}_+ with barycenter S_0 such that the set $\mathcal{M}(\nu_1, \dots, \nu_T)$ of martingale measures π satisfying $S_t(\pi) = \nu_t$ is non-empty. Let $\Phi : \mathbb{R}_+^T \rightarrow \mathbb{R}$ be u.s.c. and linearly bounded from above. Then*

$$\begin{aligned} p^M(\Phi) &:= \sup_{\pi \in \mathcal{M}(\nu_1, \dots, \nu_T)} \left\{ \int_{\mathbb{R}_+^T} \Phi(x) d\pi(x) \right\} \\ &= \inf \left\{ \sum_{t=1}^T \int_{\mathbb{R}_+} \varphi_t d\nu_t : \varphi_t \in L^1(\nu_t), \exists \Delta \in \mathcal{H} \text{ s.t. } \sum_{t=1}^T \varphi_t(x_t) + (\Delta \cdot x)_T \geq \Phi(x) \right\} =: p^R(\Phi). \end{aligned}$$

In addition, the above supremum is a maximum.

For us this follows precisely in the same way as Corollary 4.3, by including the options $\{\pm \tilde{\psi}_{k,t}, k \in \mathbb{R}_+, t = 1, \dots, T-1\}$ among the $(\varphi_i)_{i \in I}$, where

$$\tilde{\psi}_{k,t}(x) := (x_t - k)_+ - \int_k^\infty (y - k) d\mu_t(y).$$

5. CONNECTION WITH MARTINGALE-INEQUALITIES

In this section we illustrate how the Super-Replication Theorem 1.4 connects to the field of martingale inequalities. We will concentrate on the particular case of the Doob- L^1 inequality. In its sharp version obtained by Gilat [Gil86] it asserts that for every non-negative martingale $S = (S_t)_{t=0}^T$ starting at $S_0 = 1$ we have

$$\mathbb{E}[\bar{S}_T] \leq \frac{e}{e-1} \left[\mathbb{E}[S_T \log(S_T)] + 1 \right], \quad (5.1)$$

where \bar{S}_T is the supremum of S up to time T .

Having Theorem 1.4 in mind, it is natural to ask whether there exists a path-wise hedging inequality associated to it.

Claim. Fix $C \geq 0$. For every $\varepsilon > 0$ there exist $a \geq 0$ and Δ such that

$$\bar{x}_T \leq a(x_T \log(x_T) - C) + \frac{e}{e-1}(C+1) + \varepsilon + (\Delta \cdot x)_T \quad (5.2)$$

for all $x_1, \dots, x_T \in \mathbb{R}_+$.

Proof. Fix C and ε . To establish a connection with the robust Super-Replication Theorem, we let $x_0 := 1$ and interpret $\Phi(x_1, \dots, x_T) := \bar{x}_T = \max(x_0, \dots, x_T)$ and $\varphi(x_T) := x_T \log(x_T)$ as financial derivatives, where φ can be bought at price C on the market. Our task is then to determine a reasonable upper bound for the price of Φ . (Note that from $C \geq 0$ it follows that the set of admissible martingale measures is non-empty as witnessed by the a.s. constant process $S \equiv 1$.) By (5.1) we have

$$\sup \left\{ \int_{\mathbb{R}_+^T} \bar{x}_T d\pi : \pi \in \mathcal{M}, \int_{\mathbb{R}_+^T} x_T \log(x_T) d\pi \leq C \right\} \leq \frac{e}{e-1}(C+1), \quad (5.3)$$

where \mathcal{M} is the set of all martingale measures. Applying Theorem 1.4 to Φ and $\varphi_0 := \varphi - C$ we thus obtain that \bar{S}_T can be super-replicated path-wise using an initial endowment of at most $\frac{e}{e-1}(C+1) + \varepsilon$. This is precisely what is asserted in (5.2). \square

These considerations provided the motivation to search for an explicit super-replication strategy for $\Phi(x) = \bar{x}_T$ (see [ABP⁺12]). Indeed (5.2) holds (independent of C) for the particular choices $a = \frac{e}{e-1}$, $\varepsilon = 0$ and $\Delta_t(x_1, \dots, x_t) = -\log(\bar{x}_t) \cdot x$, where it corresponds to

$$\bar{x}_T \leq \frac{e}{e-1} (x_T \log(x_T) + 1) - (\log(\bar{x}_t) \cdot x)_T. \quad (5.4)$$

Let us stress that (5.4) is simply an inequality for non-negative numbers x_1, \dots, x_T . Its verification, using convexity of $x \mapsto x \log(x)$, is entirely elementary ([ABP⁺12, Proposition 2.1]). An application of (5.4) is that it implies Doob's L^1 -inequality:

Proof of (5.1). Apply (5.4) to the paths of $(S_n)_{n=0}^T$ and take expectation to obtain

$$\begin{aligned} \mathbb{E}[\bar{S}_T] &\leq \frac{e}{e-1} [\mathbb{E}[S_T \log(S_T)] + 1] - \mathbb{E}[(\log(\bar{S}_t) \cdot S)_T] \\ &= \frac{e}{e-1} [\mathbb{E}[S_T \log(S_T)] + 1]. \end{aligned} \quad \square$$

We emphasize that by Theorem 1.4 one knows *a priori* that a path-wise hedging strategy exists and hence that the Doob L^1 -inequality can be proved *in this way*. In particular one expects that the same strategy of proof can be applied to a variety of other inequalities.

REFERENCES

- [ABP⁺12] B. Acciaio, M. Beiglböck, F. Penkner, W. Schachermayer, and J. Temme. A trajectorial interpretation of Doob's martingale inequalities. *Ann. Appl. Probab.*, 2012. To appear.

- [BHLP12] M. Beiglböck, P. Henry-Labordère, and F. Penkner. Model-independent Bounds for Option Prices: A Mass Transport Approach. *Finance Stoch.*, to appear, 2012.
- [BHR01] H. Brown, D. Hobson, and L.C.G. Rogers. Robust hedging of barrier options. *Math. Finance*, 11(3):285–314, 2001.
- [BJ12] M. Beiglböck and N. Juillet. On a problem of optimal transport under marginal martingale constraints. *ArXiv e-prints*, pages 1–51, 2012.
- [BL78] D.T. Breeden and R.H. Litzenberger. Prices of State-Contingent Claims Implicit in Option Prices. *The Journal of Business*, 51(4):621–651, oct 1978.
- [Bue06] H. Buehler. Expensive martingales. *Quantitative Finance*, 6(3):207–218, 2006.
- [Cam10] L. Campi. Super-hedging. In R. Cont, editor, *Encyclopedia of Quantitative Finance*. Wiley, 2010.
- [CHO08] A. M. G. Cox, David Hobson, and Jan Oblój. Pathwise inequalities for local time: applications to Skorokhod embeddings and optimal stopping. *Ann. Appl. Probab.*, 18(5):1870–1896, 2008.
- [CM05] P. Carr and D.B. Madan. A note on sufficient conditions for no arbitrage. *Finance Research Letters*, 2(3):125–130, 2005.
- [CO11a] A.M.G. Cox and J. Oblój. Robust hedging of double touch barrier options. *SIAM J. Financial Math.*, 2:141–182, 2011.
- [CO11b] A.M.G. Cox and J. Oblój. Robust pricing and hedging of double no-touch options. *Finance Stoch.*, 15(3):573–605, 2011.
- [Cou04] L. Cousot. Necessary and Sufficient Conditions for No Static Arbitrage among European Calls. *Courant Institute, New York University*, 2004.
- [Cou07] L. Cousot. Conditions on option prices for absence of arbitrage and exact calibration. *Journal of Banking & Finance*, 31(11):3377–3397, 2007. Risk Management and Quantitative Approaches in Finance.
- [CW12] A. M. G. Cox and J. Wang. Root’s Barrier: Construction, Optimality and Applications to Variance Options. *Ann. Appl. Probab.*, to appear, 2012.
- [DH07] M.H.A. Davis and D. Hobson. The range of traded option prices. *Math. Finance*, 17(1):1–14, 2007.
- [DM04] S. Debaris and C. Martini. Superhedging strategies and balayage in discrete time. In *Seminar on Stochastic Analysis, Random Fields and Applications IV*, volume 58 of *Progr. Probab.*, pages 205–219. Birkhäuser, Basel, 2004.
- [DOR12] M.H.A. Davis, J. Oblój, and V. Raval. Arbitrage Bounds for Prices of Options on Realized Variance. *Math. Finance*, to appear, 2012.
- [DS12] Y. Dolinsky and M. H. Soner. Robust Hedging and Martingale Optimal Transport in Continuous Time. *ArXiv e-prints*, 2012.
- [DS13] Y. Dolinsky and M. H. Soner. Robust Hedging under Proportional Transaction Costs. *preprint*, 2013.
- [GHLT11] A. Galichon, P. Henry-Labordère, and N. Touzi. A Stochastic Control Approach to No-Arbitrage Bounds Given Marginals, with an Application to Lookback Options. *SSRN eLibrary*, 2011.
- [Gil86] D. Gilat. The best bound in the $L \log L$ inequality of Hardy and Littlewood and its martingale counterpart. *Proc. Amer. Math. Soc.*, 97(3):429–436, 1986.
- [HK12] D. Hobson and M. Klimmek. Model independent hedging strategies for variance swaps. *Finance and Stochastics*, 16(4):611–649, oct 2012.
- [HLOST12] P. Henry-Labordère, J. Obloj, P. Spoida, and N. Touzi. Maximum Maximum of Martingales Given Marginals. *SSRN eLibrary*, 2012.
- [HN12] D. Hobson and A. Neuberger. Robust bounds for forward start options. *Mathematical Finance*, 22(1):31–56, dec 2012.
- [Hob98] D. Hobson. Robust hedging of the lookback option. *Finance and Stochastics*, 2:329–347, 1998. 10.1007/s007800050044.
- [Hob11] D. Hobson. The Skorokhod embedding problem and model-independent bounds for option prices. In *Paris-Princeton Lectures on Mathematical Finance 2010*, volume 2003 of *Lecture Notes in Math.*, pages 267–318. Springer, Berlin, 2011.
- [HP02] D. Hobson and J.L. Pedersen. The minimum maximum of a continuous martingale with given initial and terminal laws. *Ann. Probab.*, 30(2):978–999, 2002.
- [MY02] D.B. Madan and M. Yor. Making Markov martingales meet marginals: with explicit constructions. *Bernoulli*, 8(4):509–536, 2002.
- [Nut13] M. Nutz. Superreplication under Model Uncertainty in Discrete Time. *ArXiv e-prints*, pages 1–14, 2013.
- [Obł04] J. Oblój. The Skorokhod embedding problem and its offspring. *Probab. Surv.*, 1:321–390, 2004.
- [Rie11] F. Riedel. Finance Without Probabilistic Prior Assumptions. *ArXiv e-prints*, July 2011.
- [Rog93] L. C. G. Rogers. The joint law of the maximum and terminal value of a martingale. *Probab. Theory Related Fields*, 95(4):451–466, 1993.
- [Sch10] W. Schachermayer. The Fundamental Theorem of Asset Pricing. In R. Cont, editor, *Encyclopedia of Quantitative Finance*, volume 2, pages 792–801. Wiley, 2010.
- [TT11] X. Tan and N. Touzi. Optimal Transportation under Controlled Stochastic Dynamics. *Ann. Probab.*, to appear, 2011.